

Myelinated nerve fiber layer with visual field defect

Category(ies): Pediatrics, Retina

Contributor: [Luke Lenci, MD](#)

Photographer: Samantha Jacobs



This is an example of a prominent myelinated nerve fiber layer. These can have corresponding visual field defects as seen here on Goldmann perimetry with depression of the I4e, I3e, and I2e isopters in the region of the myelination. This is an unusual case as it is located along the horizontal raphe; typically, the myelin is peripapillary in location.



[Enlarge](#)

[Download](#)




Image Permissions:



Ophthalmic Atlas Images by EyeRounds.org, The University of Iowa are licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License.

University of Iowa
Roy J. and Lucille A. Carver College
of Medicine
Department of Ophthalmology and
Visual Sciences
200 Hawkins Drive
Iowa City, IA 52242

[Support Us](#)

Copyright © 2019 The University of
Iowa. All Rights Reserved
[Report an issue with this page](#)
[Web Privacy Policy](#) |
[Nondiscrimination Statement](#)

[Cataract Surgery for Greenhorns](#)
[EyeTransillumination](#)
[Gonioscopy.org](#)
[Iowa Glaucoma Curriculum](#)
[Iowa Wet Lab](#)
[Patient Information](#)
[Stone Rounds](#)
[The Best Hits Bookshelf](#)

Media

Follow



[Receive notification of new cases,
sign up here](#)
[Contact Us](#)
[Submit a Suggestion](#)