

Conjunctival squamous papilloma


Category(ies): Cornea, External Disease

Contributor: [Jesse Vislisel, MD](#)

Photographer: Cindy Montague, CRA



Squamous papillomas can be pedunculated or sessile as shown in this photograph. The sessile variety has a flat base and is found most commonly along the limbus. They have a glistening surface, often with red dots or "corkscrew" blood vessels, creating a strawberry-like appearance. The lesions may encroach onto the cornea and must be monitored closely for signs of dysplastic change or carcinomatous growth.



[Enlarge](#) [Download](#)

Image Permissions:



Address

University of Iowa
Roy J. and Lucille A. Carver College
of Medicine
Department of Ophthalmology and
Visual Sciences
200 Hawkins Drive
Iowa City, IA 52242

[Support Us](#)

Legal

Copyright © 2019 The University of
Iowa. All Rights Reserved
[Report an issue with this page](#)
[Web Privacy Policy](#) |
[Nondiscrimination Statement](#)

Related Links

[Cataract Surgery for Greenhorns](#)
[EyeTransillumination](#)
[Gonioscopy.org](#)
[Iowa Glaucoma Curriculum](#)
[Iowa Wet Lab](#)
[Patient Information](#)
[Stone Rounds](#)
[The Best Hits Bookshelf](#)

EyeRounds Social Media

Follow



Receive notification of new cases,
[sign up here](#)
[Contact Us](#)
[Submit a Suggestion](#)