

## Lisch nodules


**Category(ies):** [Glaucoma](#), [Iris](#), [Pediatrics](#)

**Contributor:** [Jesse Vislisel, MD](#)

**Photographer:** Ed Heffron, Brice Critser, CRA




Lisch nodules are melanocytic hamartomas of the iris, often associated with neurofibromatosis (NF) I. They are usually elevated and tan in appearance. Their incidence in NF1 increases with age and their prevalence raises by about 10% per year of life, up to age 9. These photographs show the various appearances of these nodules in 3 different patients with NF1.



Photos of patient by Brice Critser, CRA

[Enlarge](#) [Download](#)




Photos of patient by Brice Critser, CRA

[Enlarge](#)

[Download](#)





Photos of patient 3 by Ed Heffron

[Enlarge](#)

[Download](#)





Photos of patient 3 by Ed Heffron

[Enlarge](#)

[Download](#)

### Image Permissions:



Ophthalmic Atlas Images by [EyeRounds.org](#), [The University of Iowa](#) are licensed under a [Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License](#).

### Address

University of Iowa  
Roy J. and Lucille A. Carver College  
of Medicine  
Department of Ophthalmology and  
Visual Sciences  
200 Hawkins Drive  
Iowa City, IA 52242

[Support Us](#)

### Legal

Copyright © 2019 The University of  
Iowa. All Rights Reserved  
[Report an issue with this page](#)  
[Web Privacy Policy](#) |  
[Nondiscrimination Statement](#)

### Related Links

[Cataract Surgery for Greenhorns](#)  
[EyeTransillumination](#)  
[Gonioscopy.org](#)  
[Iowa Glaucoma Curriculum](#)  
[Iowa Wet Lab](#)  
[Patient Information](#)  
[Stone Rounds](#)  
[The Best Hits Bookshelf](#)

### EyeRounds Social Media

Follow



[Receive notification of new cases,  
sign up here](#)  
[Contact Us](#)  
[Submit a Suggestion](#)